




The Open Biology Journal

Content list available at: <https://openbiologyjournal.com/>



RESEARCH ARTICLE

Different Effects of Fluoxetine and Paroxetine Combined with Vitamin D₃ in Ovariectomized Rats Exposed to Unpredictable Stress

Julia Fedotova^{1,2,*}

¹Laboratory of Neuroendocrinology, I.P. Pavlov Institute of Physiology RASci., 6 Emb. Makarova, St. Petersburg 199034, Russia

²The International Research Centre "Biotechnologies of the Third Millennium", ITMO University, 9 Lomonosova Str., St. Petersburg 191002, Russia

Abstract:

Background:

Vitamin D₃ (VD₃) is involved in the pathophysiological mechanisms of affective-related disorders and controls the functional activity of various hormonal systems. The complex interaction between estrogen and VD₃ creates a neurobiological basis for their participation in similar behavioral disorders.

Objectives:

This study aimed to evaluate whether VD₃ (5.0 mg/kg, s.c.) facilitates the antidepressant-like action of fluoxetine (10.0 mg/kg, i.p.) or paroxetine (10.0 mg/kg, i.p.) by enhancing the antidepressant-like activity of these drugs in adult long-term Ovariectomized (OVX) rats subjected to Chronic Unpredictable Mild Stress (CUMS) protocol for 6 weeks.

Methods:

Sucrose Preference (SPT) and Forced Swim (FST) tests were performed to evaluate the anhedonia state and depressive symptoms, respectively. The Open-Field Test (OFT) was carried out to measure locomotor activity as well as grooming behavior produced by CUMS in long-term OVX rats. Corticosterone (CS)/estradiol (E₂) in the serum was tested by rat ELISA kits. NF-kB, 5-HT/5-HIAA, and pro-inflammatory cytokine levels in the hippocampus were also examined by rat ELISA kits.

Results:

The results of this study suggest that combined treatment with fluoxetine (10.0 mg/kg, i.p.) or paroxetine (10.0 mg/kg, i.p.) along with VD₃ (5.0 mg/kg, s.c.) produces distinct effects on the depression-like behavior in long-term OVX/CUMS rats. Co-administration of fluoxetine (10.0 mg/kg, i.p.) with VD₃ did not facilitate the antidepressant-like effects of fluoxetine in the long-term OVX rats with CUMS. On the other hand, co-treatment with paroxetine with VD₃ resulted in faster and marked antianhedonic- and antidepressant-like effects in long-term OVX rats with CUMS as compared to treatment with paroxetine alone. The co-administration of paroxetine and VD₃ attenuates stress-induced modifications of CS/E₂ levels in the serum, as well as proinflammatory cytokine/NF-kB/5-HT levels in the hippocampus of long-term OVX rats exposed to CUMS.

Conclusion:

Supplementation of VD₃ (5.0 mg/kg, s.c.) to paroxetine (10.0 mg/kg, i.p.) facilitates antianhedonic- and antidepressant-like effects of paroxetine in adult long-term OVX rats exposed to CUMS.

Keywords: Vitamin D₃, Fluoxetine, Paroxetine, Long-term ovariectomy, Chronic unpredictable mild stress, 5-HT, Pro-inflammatory cytokines.

Article History

Received: May 10, 2020

Revised: August 20, 2020

Accepted: September 17, 2020

1. INTRODUCTION

Estrogen deficiency during menopause can lead to pro-

found endocrine disorders and neuropsychiatric symptoms, such as memory impairment, insomnia, anxiety, and depression [1 - 3]. These disorders have an adverse impact on the quality of life [4]. Approximately 36% of the menopausal women report experiencing depressive symptoms [5].

Despite numerous neuropharmacological schedules and

* Address correspondence to this author at the Laboratory of Neuroendocrinology, I.P. Pavlov Institute of Physiology, Russian Academy of Sciences, 6 Emb. Makarova, St. Petersburg 199034, Russia; Tel: +7 911 287 92 73, Fax: +7 812 328 05 01; E-mail: julia.fedotova@mail.ru

psychological rehabilitation approaches to treat mood disorders, only moderate improvements have been registered in the treatment of affective-related disorders. More importantly, the possibility of recurrence is increased especially in women during estrogen deficiency [6, 7]. The standard treatment of mood disorders with antidepressants highlights the restricted effectiveness of the current approach [8 - 11].

Among the different classes of antidepressants, SSRIs are the most extensively used pharmacological drugs to treat affective-related disorders in perimenopausal and postmenopausal women [3, 9, 11]. However, the currently used antidepressants are not completely effective and have many adverse effects [6 - 8]. Moreover, the therapeutic response to SSRIs in perimenopausal women might be influenced by modifications in the concentrations of female gonadal hormones that can restrict their application in such cases [12, 13]. The absence or delayed response to treatment with antidepressants is a major problem in the therapy of depression [8 - 11].

The monoamines hypothesis of depression suggests that depression arises from the decreasing release of monoamines, especially serotonin (5-HT) in the brain [14]. Another important factor behind the onset of depression is inflammation in the brain [15 - 19]. Depressed patients have been documented to have higher levels of serum pro-inflammatory cytokines, such as interleukin (IL)-1 β , IL-6, and tumor necrosis factor-alpha (TNF- α) [19, 20]. Moreover, increased inflammatory mediator levels may act as predictive biomarkers to resist the pharmacotherapy of depression [21]. Furthermore, nuclear factor-kappa B (NF- κ B) is one of the most important pro-inflammatory transcription factors that is involved in the pathophysiology of numerous peripheral and central inflammatory impairments [22, 23]. NF- κ B is activated by stress and might mediate molecular outcomes to stress factors and trigger mood deteriorations [24, 25]. Findings from clinical studies indicated that women in the menopausal period have a significant imbalance in the production of inflammatory molecules as well as decreased serotonergic activity [26, 27].

Vitamin D (VD) deficiency is one of the many factors that lead to the onset of mood disturbances [28]. VD is synthesized in the skin and can be also found in food [29, 30]. A strong relationship between VD deficiency and profound affective-related state, as well as between VD deficiency and cognitive function have been reported in several studies [31 - 34].

Estrogen imbalance can lead to symptoms of VD deficiency since estrogen induces the increased activity of the enzyme system causing VD action [35]. Some clinical studies showed that mild depressive symptoms are more frequently registered in women with VD insufficiency, and even more so in women with VD deficiency [36, 37]. In addition, VD₃ controls the functional activity of the 5-HT system in the brain [38]. VD₃ might transcriptionally activate the tryptophan hydroxylase-2 gene, resulting in increased conversion of tryptophan to 5-HT in the brain [38, 39]. Moreover, VD₃ modulates neuroinflammation in the brain and decreases proinflammatory cytokine levels in the central and peripheral nervous system [40, 41].

Altogether, the above-mentioned findings suggest that mood disturbances in menopausal women may involve complex deterioration in estrogen and VD₃ levels, neuroinflammation, as well as abnormal 5-HT levels in the brain.

In our previous studies, we assessed the antidepressant-like effects of VD₃ at different doses (1.0, 2.5, and 5.0 mg/kg, s.c.) on a model of depression induced by Chronic Unpredictable Mild Stress (CUMS) for 28 days in adult ovariectomized (OVX) rats [42]. The results of our study demonstrated that VD₃ (5.0 mg/kg, s.c.) considerably reversed the depression-like state in the Forced Swimming Test (FST) in long-term OVX rats compared to OVX rats with CUMS [42]. Therefore, we designed the current study taking into account the antidepressant-like activity of VD₃ in the affective-related profile of the long-term OVX females.

This study aimed to evaluate whether VD₃ facilitates the antidepressant-like action of fluoxetine or paroxetine by enhancing the antidepressant-like activity of these SSRIs in female rats with long-term deficiency of estrogen. Fluoxetine and paroxetine are well-known SSRIs that are frequently prescribed in the treatment of depression [43, 44]. Moreover, paroxetine is also suspected to modulate the female reproductive system and estradiol levels in postmenopausal women [45, 46]. The effects of co-administration of fluoxetine or paroxetine with VD₃ were evaluated by the sucrose preference test, forced swimming test, and open-field test in adult female rats with long-term deficiency of estrogen exposed to the CUMS. The effects of co-administration with VD₃ and fluoxetine/paroxetine on changes in NF- κ B signaling, pro-inflammatory cytokines, and 5-HT levels in the hippocampus were evaluated in the adult female rats with long-term deficiency of estrogen exposed to the CUMS.

2. MATERIALS AND METHODS

2.1. Animals

Female Wistar rats (190 \pm 20 g) were purchased from the Animal Rat Center at the Rappolovo Laboratory Animal Factory (St. Petersburg, Russia). A total number of 56 female rats at the age of 3 months at the beginning of the experiment were divided into different groups. Rats were housed singly at a controlled temperature (22 \pm 1 $^{\circ}$ C), humidity (50 \pm 10%), and light conditions with *ad libitum* access to food and water. The light was programmed on a 12h light/dark cycle (lights on at 7:00 AM). The experimental females were assigned to stay at the animal room for 1 week before their use in the experiment. All procedures with animals were carried out in compliance with national and international laws and policies. Experimental rats were decapitated with anesthesia for brain sampling and every effort was made to minimize the pain experienced by the animals. In all stress procedures, the researchers treated the rats with care and were sensitive to their emotional state.

2.2. Surgery Protocol

Ovariectomy was performed as previously described [42]. The rats were anesthetized with narcosis that comprised of a mixture of 10 mg/kg xylazine and 70 mg/kg ketamine for bilateral removal of the ovaries. Ovariectomy was performed

using 2 typical cuts in a lateral position. Under anesthesia, a longitudinal incision was made along the dorsal midline. The muscle layer was drawn from the abdomen through the incision with sterilized hemostatic forceps. The oviducts were clamped with hemostatic forceps and the ovaries were removed. The muscle layer was then closed. The efficiency of the surgery was examined by routine vaginal inspection. The sham ovariectomy was performed in an identical way but without the removal of the female endocrine glands. After the ovariectomy and sham operation, the experimental rats were maintained in their home cage and were allowed to recover for a period of 3 months as a post-ovariectomy period. During the post-surgery interval, experimental groups of females admitted were given free access to food and water. After 3 months of the post-ovariectomy period, rats were assigned randomly to the experiment for chronic stress procedure, except for the control sham-operated (SHAM) rats.

2.3. Chronic Unpredictable Mild Stress Trial

The Chronic Unpredictable Mild Stress (CUMS) protocol was performed as reported previously [47, 48]. Rats in CUMS groups were submitted to different stressful stimuli for 14 days, followed by 28 days (from third to the sixth week) of VD₃, fluoxetine, paroxetine, VD₃, and fluoxetine, and VD₃ and paroxetine treatment for which the stress protocol continued. The following stressors were used: water deprivation (24 h), food deprivation (24 h), wet bedding (24 h), tilted cage at 45° (24 h), a reversal of the light/dark cycle (24 h), swimming in cold water (4 °C, 5 min), swimming in warm water (45 °C, 5 min), and clippedtail (1 min, 1 cm from the end of the tail). The rats were exposed to one of the stressors on a given day. All stress factors were applied individually and continuously throughout the day. To avoid habituation, the same stressor was not applied consecutively for two days to guarantee that the animal does not predict the occurrence of any stress trigger. CUMS protocol was similar every time to maintain reproducibility. During the stress trial, the rats were placed into the experimental room to undergo the stress procedure and were returned to their single cages afterwards. The stress procedure did not involve any food or water deprivation in the present study. The females after sham-operation (control rats) were placed in a separate cage and vivarium section without any contact with the CUMS groups of animals for 6 weeks. The rats were generally undisturbed and well maintained (provided with food and routine cage cleaning). All experimental females were supervised daily by the veterinary care staff during sub-routine maintenance. The rats were examined in a special animal room using a non-invasive observational assessment procedure that yielded information regarding the health of each animal. The assessment consisted of several measurements, such as body condition, appearance, breathing, hydration status, posture, mobility, muscle tone, as well as any presence of defects in the bones, genitals, and abdomen. No rats were injured or unhealthy during the experimental protocol.

2.4. Drug Treatments

The pharmacological preparations used in the current study were fluoxetine hydrochloride, paroxetine hydrochloride, and cholecalciferol as VD₃. All of the drugs were obtained from Sigma Chemical Co. (St. Louis, MO, USA). Fluoxetine hydrochloride and paroxetine hydrochloride were given Intraperitoneally (IP). VD₃ was administered Subcutaneously (SC) in a volume of 0.1 mL/rat, respectively, for 21 days during the CUMS procedure: 1 hour before the daily stressor action, and again during the period of the behavioral testing. Behavioral evaluations were recorded 60 minutes after the final drug treatment. Fluoxetine hydrochloride and paroxetine hydrochloride were dissolved in sterile 0.9% saline daily. Cholecalciferol was dissolved in a solvent of 95% ethanol, and then aliquoted and maintained at -80°C. Cholecalciferol solution for SC injection was freshly prepared. VD₃ was diluted in physiological saline water, resulting in a solvent of VD₃ with 1% ethanol.

2.5. Groups of Animals

All animals were randomly assigned to a total of 8 experimental groups ($n = 7$ in each):

- (1) sham-operated (SHAM) rats without the CUMS model treated with solvent (control).
- (2) SHAM rats exposed to CUMS treated with solvent.
- (3) long-term OVX rats exposed to CUMS treated with solvent.
- (4) long-term OVX rats exposed to CUMS treated with VD₃ (5.0 mg/kg/day, SC).
- (5) long-term OVX rats exposed to CUMS treated with fluoxetine (10.0 mg/kg/day, IP).
- (6) long-term OVX rats exposed to CUMS treated with fluoxetine and VD₃.
- (7) long-term OVX rats exposed to CUMS treated with paroxetine (10.0 mg/kg/day, i.p.).
- (8) long-term OVX rats exposed to CUMS treated with paroxetine and VD₃.

In preliminary our works, no significant differences were found between SHAM/OVX rats treated with physiological saline (0.9% saline) as the solvent for fluoxetine, and SHAM/OVX females treated with a special solvent for VD₃ in behavioral trials (data are not shown). Therefore, physiological saline was used as the solvent for SHAM/OVX female rats. The doses of VD₃ were based on our previous studies on the behavioral effects of VD₃ on depression-like behavior of non-stressed long-term OVX female rats [42]. Doses of fluoxetine and paroxetine were chosen according to reported experimental data [49, 50]. All behavioral measurements were recorded 60 minutes after the last drug administration. To minimize animal suffering, all groups of rats were euthanized by pentobarbital overdose after all behavioral trials. The timeline of this experimental study is presented in Fig. (1).




Fig. (1). Scheme of chronic treatment with SSRIs and VD₃. Female rats were divided into 8 groups: 1 – control SHAM; 2 – SHAM + CUMS + solvent; 3 – OVX + CUMS + solvent; 4 – OVX + CUMS + VD₃ (5.0 mg/kg, s.c.); 5 – OVX + CUMS + fluoxetine (10.0 mg/kg, i.p.); 6 – OVX rats + CUMS + fluoxetine (10.0 mg/kg, i.p.) + VD₃ (5.0 mg/kg, s.c.); 7 – OVX + CUMS + paroxetine (10.0 mg/kg, i.p.); 8 – OVX rats + CUMS + paroxetine (10.0 mg/kg, i.p.) + VD₃ (5.0 mg/kg, s.c.).

2.6. Body Weight

The body weight of all experimental groups was evaluated every week during the whole CUMS protocol.

2.7. Sucrose Preference Test

Sucrose Preference Test (SPT) was conducted every week during the 6 weeks of the CUMS protocol [44]. The SPT was performed as reported previously [42]. All groups of experimental rats were exposed to a 24-hour deprivation of food and water. After twenty-four hours, the subjects had 1-hour access to a bottle of water and a bottle of sucrose solution of 200 ml each. The measurement of sucrose preference was calculated using the following formula: % sucrose preference = sucrose intake/sucrose intake plus water intake × 100%.

2.8. Forced Swimming Test

The Forced Swimming Test (FST) was conducted as described previously [42]. During the FST test, the experimental female rats were put into the FST cylinders (height 60 cm and diameter 20 cm) that were filled with 23-25 °C water up till a depth of 30 cm. The immobility time, swimming time, and climbing time were recorded for 5 minutes using a video camera that was placed above the apparatus.

2.9. Open Field Test

The Open Field Test (OFT) was carried out as previously performed [42]. In brief, the animals were set in the center square of the OFT apparatus and observed for 300 seconds. The number of total crossings, the number of crossings in the central squares, the number of rearings, the time spent in the central squares, and grooming were measured for 300 seconds in the OFT using a video camera. After each animal's trial, the OFT apparatus was cleaned thoroughly with 75% alcohol to prevent smell interference between trials.

2.10. Serum Corticosterone and Estradiol Measurements

Following behavioral tests, all rats were decapitated under deep anesthesia (10 mg/kg xylazine and 70 mg/kg ketamine) for brain sampling. 5 mL samples of blood were collected from the animals and were centrifuged at 4000 ×g for 15 minutes at 4°C. The brain tissue samples needed for ELISA assay were stored in a -80°C refrigerator for later use. The serum samples were used to estimate the Corticosterone (CS) and Estradiol (E₂) levels using commercially available rat ELISA kits (Cusabio Biotech Co., Ltd, Wuhan, China) according to the manufacturer's directions. The sensitivity and detection range of the CS were 0.1 ng/mL and 0.2-40 ng/mL, respectively, and 40.0 pg/mL and 40.0-1500 pg/mL for E₂, respectively.

2.11. Hippocampal NF-kB/p50/p65, TNF-α, IL-1β, IL-6 Levels Determination

The cold lysis extraction buffer (0.2% sodium deoxycholate, 0.5% Triton X-100, 1% NP-40, 50 mM Tris-HCl pH 7.4, 1 mM phenylmethylsulfonyl fluoride, 1 mM N-ethylmaleimide, and 2.5 mM phenanthroline) was used for the dissection and homogenization of the hippocampi of the experimental groups of rats [51]. The brain samples with cold lysis buffer were then sonicated for 15 seconds. After that, the hippocampi were centrifuged at 12,000 ×g for 15 minutes at 4°C. To normalize the hippocampi supernatants to the total protein, the Bradford method was applied [52]. The protein normalized supernatants of the hippocampus were stored at -80°C until assays were conducted.

Homogenates of the hippocampus in all experimental groups were utilized for evaluation of the levels using rat nuclear factor-kB/p50/p65 (NF-kB/p65/p50), tumor necrosis factor-α (TNF-α), interleukin-1-β (IL-1β), interleukin-6 (IL-6) ELISA kits (Cusabio Biotech Co., Ltd, Wuhan, China) in accordance with the manufacturer's recommendations. The NF-kB/p50/p65, TNF-α, IL-1β, IL-6 levels were detected using an MC Thermo Fisher Scientific reader (Thermo Fisher Sci-

entific Inc., Helsinki, Finland) at 450 nm. The NF- κ B/p65/p50 content is presented as pg/mg of tissue. The sensitivity and detection range of the NF- κ B/p65 rat ELISA kit were 10.0 pg/ml and 62.5 pg/ml-2000 pg/ml, respectively. The sensitivity and detection range of the NF- κ B/p50 rat ELISA kit were 0.119 ng/ml and 0.312 ng/ml-20 ng/ml. The TNF- α , IL-1 β , IL-6 levels were expressed as pg/mL. The sensitivity and detection range of the TNF- α rat ELISA kit were 1.56 pg/mL and 6.25-400 pg/mL, respectively; for IL-1 β rat ELISA kit - 1.56 pg/mL and 6.25-4000 pg/mL, respectively; for IL-6 rat ELISA kit - 0.078 pg/mL and 0.312-20 pg/mL, respectively. The determination showed no significant cross-reference with other biological factors. The duplication of all samples was performed for the ELISA assay.

2.12. Hippocampal 5-HT and 5-HIAA Levels Measurement

The 5-HT/5-HIAA levels in the hippocampus were examined using High-Performance Liquid Chromatography (HPLC) as previously reported [53]. All hippocampal samples were homogenized in a 0.1 mol/L solution of HClO₄ along with 0.02% Na₂S₂O₂ (15 μ L of solution for each mg of tissue) and Dihydroxybenzylamine (DHBA, 146.5 ng/mL, internal standard). These homogenates were centrifuged for 40 minutes at 11,000 \times g at 4°C. The isocratic system (Shimadzu LC-10AD, Kyoto, Japan) in HPLC equipment was supplied by a Spheri-5 RP-18 5 μ m column (220 \times 4.6 mm) and a 20 μ L injection loop. The electrochemical detection was 0.75 V. The mobile phase consisted of 0.06% heptane sulphonic acid and phosphate/citrate pH 2.64, 0.02 mol/L, 0.12 mmol/L ethylene diamine tetraacetic acid, containing 10% methanol. The flow rate was 1 mL/min. 5-HT and its metabolite levels were expressed as mean \pm SD ng/mg tissue. A solution containing 2 mL of sodium tetraborate 0.1 mol/L and 1 mL of stock solution was prepared before the assay of hippocampi. To finalize the precolumn derivatization, we used 100 μ L of this solution with 50 μ L of the sample for 2 minutes before each injection [53]. For the mobile phase, sodium phosphate 0.05 mol/L (pH 5.95) with 11.5% methanol were applied. HPLC system had a flow rate of 3.5 mL/min. The emission was detected at 460 nm, and excitation was detected at 348 nm. Standards of 5-HT and 5-HIAA were validated before determination. We did not find any additional peaks except 5-HT or 5-HIAA. The retention time was simultaneously detected for -HT and 5-HIAA.

2.13. Statistical Analysis

All experimental values are presented as mean \pm Standard Error of the Mean (SEM). The differences in body weight and sucrose preference consumption among all experimental groups were analyzed using the two-way analysis of variance (two-way ANOVA) test with repeated measurements between subject factors for the duration and drug treatments. The other behavioral and biochemical evaluations between all experimental groups were analyzed using one-way ANOVA without repeated measurements followed by the *post hoc* Tukey test for multiple comparisons. The statistical analysis was carried out using the Statistics Package for SPSS, version 18.0 (SPSS Inc., USA). The probability level of $P < 0.05$ was considered to be statistically significant.

3. RESULTS

3.1. Combined Treatment (SSRIs with VD₃) Increases Body Weight in Long-term OVX Rats Exposed to CUMS

Before the CUMS protocol, we noted no significant difference in body weight among all experimental groups (Fig. 2). After 14 days of the CUMS trials, the ANOVA test indicated a significant effect of treatment ($F(5,34) = 15.44$, $p < 0.05$), effect of duration ($F(5,34) = 11.09$, $p < 0.05$), and a statistical effect of duration \times treatment interaction ($F(5,34) = 9.08$, $p < 0.05$) on the body weight of the experimental groups. The difference in body weight among the groups ($p < 0.05$) was found using the post-hoc test. The body weight of SHAM rats subjected to CUMS gradually decreased after 2 weeks of paradigm protocol compared to non-CUMS group (post-hoc test, $P < 0.05$, (Fig. 2)). After 6 weeks of the CUMS procedure, the body weight of OVX/CUMS rats significantly reduced compared to the control non-CUMS and SHAM/CUMS groups ($P < 0.05$, (Fig. 2)). Following 28 days of different treatments, the body weight of OVX/CUMS rats treated with VD₃ (5.0 mg/kg, s.c.), fluoxetine (10.0 mg/kg, i.p.) or paroxetine (10.0 mg/kg, i.p.) was significantly higher compared to OVX/CUMS/solvent and SHAM/CUMS/solvent rats ($P < 0.05$, (Fig. 2)). The combination of fluoxetine along with VD₃ (5.0 mg/kg, s.c.) failed to influence the body weight gain of OVX/CUMS rats as compared to the OVX/CUMS/fluoxetine rats. However, the two-way ANOVA revealed a significant increase in body weight in long-term OVX rats treated with paroxetine and VD₃ after 1 week of therapy (at the 4th week of CUMS protocol) and maintained the same results in the following weeks as compared to the OVX/CUMS/solvent/paroxetine and SHAM/CUMS/solvent rats ($P < 0.05$, (Fig. 2)).

3.2. Combined Treatment (SSRIs with VD₃) Restores Sucrose Preference in Long-term OVX Rats Exposed to CUMS

The sucrose preference of long-term OVX rats subjected to CUMS and treated with SSRIs in a combination with VD₃ is presented in Fig. (3). A two-way ANOVA revealed significant differences in the sucrose preference between duration ($F(5,34) = 11.41$, $P < 0.05$), between drug treatment ($F(5,34) = 15.07$, $P < 0.05$), and an interaction between duration and treatments ($F(5,34) = 2.98$, $P < 0.05$) in the long-term OVX/CUMS rats. The post-hoc test revealed differences among the groups for sucrose preference ($P < 0.05$). We found no difference in the initial sucrose preference in all the experimental groups ($P > 0.05$, (Fig. 3)). Following CUMS trials, all experimental rats exhibited a decrease in sucrose preference in the 3rd week when compared with the control non-CUMS SHAM group ($P < 0.05$, (Fig. 3)). Moreover, the ANOVA test showed a significant effect of treatment ($F(5,12) = 11.22$, $p < 0.05$), a significant effect of duration ($F(6,22) = 21.18$, $p < 0.05$) as well as a significant effect of week \times treatment interaction ($F(25,12) = 15.14$, $p < 0.05$) on the sucrose preference in the CUMS groups. During the next 4 weeks of the CUMS trials, the SHAM/CUMS/solvent rats demonstrated a gradual decrease in sucrose preference as compared to the control non-CUMS SHAM group. The sucrose preference in long-term OVX rats from the 4th week till the end of the CUMS protocol

progressively decreased as compared to the non-CUMS/CUMS SHAM rats ($P < 0.05$, (Fig. 3)). VD_3 (5.0 mg/kg, s.c.) increased sucrose preference in the 6th week in long-term OVX rats with CUMS when compared to the OVX group subjected to CUMS and treated with the solvent ($P < 0.05$, (Fig. 3)). Treatment with fluoxetine or paroxetine, as well as fluoxetine along with VD_3 ,

steadily elevated the sucrose preference of OVX/CUMS rats in the 5th week of CUMS protocol when compared to OVX/CUMS/solvent and SHAM/CUMS/solvent rats ($P < 0.05$, (Fig. 3)). Combined administration of paroxetine and VD_3 gradually increased the sucrose preference in OVX/CUMS rats in the 4th week when compared with OVX/CUMS/ paroxetine/solvent and SHAM/CUMS/solvent rats ($P < 0.05$, (Fig. 3)).




Fig. (2). Modifications of body weight in long-term OVX rats exposed to CUMS following SSRIs alone or in a combination with VD_3 treatments. * $P < 0.05$ versus the control group, # $P < 0.05$ versus the SHAM group exposed to CUMS, \$ $P < 0.05$ versus the OVX group subjected to CUMS, \$\$ $P < 0.05$ versus the OVX group subjected to CUMS treated with fluoxetine or paroxetine, respectively. The data are presented as mean \pm SD; $n = 7$ in each group.




Fig. (3). Combined treatment with paroxetine plus VD_3 restores sucrose preference in long-term OVX rats exposed to CUMS. * $P < 0.05$ versus the control group, # $P < 0.05$ versus the SHAM group exposed to CUMS, \$ $P < 0.05$ versus the OVX group subjected to CUMS, \$\$ $P < 0.05$ versus the OVX group subjected to CUMS treated with fluoxetine or paroxetine, respectively. The data are presented as mean \pm SD; $n = 7$ in each group.




Fig. (4). Combined treatment with paroxetine plus VD₃ corrects depression-like behavior in FST of long-term OVX rats exposed to CUMS. **A** – Immobility time, sec, **B** – Swimming time, sec, **C** – Climbing time, sec. * P < 0.05 versus the control group, # – P < 0.05 versus the SHAM group exposed to CUMS, \$ – P < 0.05 versus the OVX group subjected to CUMS, \$\$ – P < 0.05 versus the OVX group subjected to CUMS treated with fluoxetine or paroxetine, respectively. The data are presented as mean ± SD; n = 7 in each group.

3.3. Combined Treatment (SSRIs with VD₃) Corrects Depression-like Behaviour in FST of Long-term OVX Rats Exposed to CUMS.

CUMS produced a significant increase in immobility time and a decrease in swimming time in the long-term OVX rats compared to the non-CUMS/SHAM/CUMS rats ($F(5,12) = 52.84$ and $F(5,12) = 68.89$, respectively, $P < 0.05$, (Fig. 4A, B).

VD₃ (5.0 mg/kg, s.c.) decreased the immobility time and increased the swimming time of OVX/CUMS rats as compared to the OVX/CUMS/solvent group ($P < 0.05$, (Fig. 4A, B). Treatment with fluoxetine and paroxetine alone significantly reduced immobility time and increased swimming time in long-term OVX/CUMS rats as compared to the OVX/CUMS/solvent and SHAM/CUMS rats ($P < 0.05$, (Fig. 4A, B).

The values of immobility time and swimming time in OVX/CUMS/fluoxetine along with VD₃ were similar to the values of OVX/CUMS/fluoxetine rats ($P < 0.05$, (Fig. 4A, B)). Paroxetine in combination with VD₃ improved immobility time and swimming time in long-term OVX rats as compared to the OVX/CUMS/paroxetine/solvent and SHAM/CUMS/ solvent rats ($P < 0.05$, Fig. (4A, B)). We found no difference in the climbing time in all the experimental groups compared to the OVX/SHAM/CUMS groups ($P > 0.05$, Fig. (4C)).


3.4. Combined Treatment (SSRIs with VD₃) Changes Behaviour in OFT of Long-term OVX Rats Exposed to CUMS

After 6 weeks, the number of total crossings, crossings in the central squares, time spent in the central squares, and rearings of SHAM/CUMS rats significantly decreased compared to the non-CUMS SHAM group ($F(5,12) = 64.43$, $P < 0.05$, (Fig. 5A-D)).

The number of total crossings, crossings in the central squares, time spent in the central squares, and rearings of long-

term OVX rats with CUMS significantly decreased compared to the non-CUMS/CUMS SHAM groups ($P < 0.05$, (Fig. 5A-D)). Administration of VD₃ enhanced the number of total crossings, crossings in the central squares, time spent in the central squares, and rearings of long-term OVX/CUMS rats compared to OVX/SHAM/CUMS groups ($P < 0.05$, (Fig. 5A-D)). Treatment with fluoxetine or paroxetine alone increased the number of total crossings, crossings in the central squares, time spent in the central squares, and rearings of long-term OVX/CUMS rats compared to the OVX/CUMS/solvent rats and SHAM/CUMS rats similarly to the OVX/CUMS rats treated with VD₃ ($P < 0.05$, (Fig. 5A-D)). The crossings in the central squares and the time spent in the central squares of long-term OVX/CUMS rats after co-administration with fluoxetine or paroxetine in a combination with VD₃ were markedly enhanced compared to OVX/CUMS/solvent/ fluoxetine/ paroxetine rats and SHAM/CUMS rats.

Following six months of CUMS protocol, we observed no statistically significant differences for grooming activities between all the experimental groups of rats in the OFT ($F(5,12) = 0.82$, $P > 0.05$, (Fig. 5E)).






Fig. (5). The behavior of long-term OVX rats exposed to CUMS after SSRIs alone or in a combination with VD₃ treatments. **A** – the number of total crossings, **B** – the number of crossings in the central squares, **C** – the number of rearings, **D** – the time spent in the central squares, **E** – grooming. * P < 0.05 versus the control group, # – P < 0.05 versus the SHAM group exposed to CUMS, \$ – P < 0.05 versus the OVX group subjected to CUMS, \$\$ – P < 0.05 versus the OVX group subjected to CUMS treated with fluoxetine or paroxetine, respectively. The data are presented as mean ± SD; n = 7 in each group.




Fig. (6). Effects of combined treatment with SSRIs plus VD_3 modulates serum corticosterone (A) and estradiol (B) levels in long-term OVX rats exposed to CUMS. * $P < 0.05$ versus the control group, # $P < 0.05$ versus the SHAM group exposed to CUMS, \$ $P < 0.05$ versus the OVX group subjected to CUMS, \$\$ $P < 0.05$ versus the OVX group subjected to CUMS treated with fluoxetine or paroxetine, respectively. The data are presented as mean \pm SD; $n = 7$ in each group.

3.5. Combined Treatment (SSRIs with VD_3) Alters Serum Corticosterone and Estradiol Levels in Long-term OVX rats Exposed to CUMS

The ELISA assay revealed that CUMS significantly increased CS concentrations in the blood of SHAM rats compared to the non-CUMS (control) female rats ($P < 0.05$, (Fig. 6A)). The serum CS levels were elevated in the long-term OVX/CUMS rats compared to the non-CUMS/SHAM/CUMS groups ($F(5,12) = 78.56$ and $F(5,12) = 56.22$, respectively, $P < 0.05$, (Fig. 6A)). Supplementation with VD_3 decreased CS and increased E_2 levels of long-term OVX/CUMS rats compared to the OVX/CUMS/solvent rats ($P < 0.05$, (Fig. 6A, B)). SSRIs significantly reduced serum CS levels in the long-term OVX rats exposed to CUMS as compared to the OVX/SHAM plus solvent rats subjected to CUMS ($P < 0.05$, (Fig. 6A)). SSRIs in a combination with VD_3 completely restored the serum CS levels in the long-term OVX rats exposed to CUMS when compared to the OVX/SHAM/fluoxetine/paroxetine rats subjected to CUMS ($P < 0.05$, (Fig. 6A)). Fluoxetine did not change serum E_2 levels in the long-term OVX rats exposed to


CUMS in comparison with the OVX/SHAM plus solvent rats subjected to CUMS ($P < 0.05$, (Fig. 6B)). Co-administration of fluoxetine with VD_3 significantly increased serum E_2 levels in long-term OVX rats as compared to the OVX/CUMS/solvent and SHAM/CUMS/solvent rats ($P < 0.05$, (Fig. 6A, B)). Paroxetine alone enhanced serum E_2 levels in long-term OVX rats when compared to OVX/CUMS/solvent and SHAM/CUMS/solvent rats ($P < 0.05$, (Fig. 6B)). The combination of paroxetine and VD_3 profoundly increased E_2 levels in long-term OVX rats when compared to OVX/CUMS/ paroxetine/solvent and SHAM/CUMS/solvent rats ($P < 0.05$, (Fig. 6B)).

3.6. Combined Treatment (SSRIs with VD_3) Modulates Hippocampal NF- κ B and Pro-inflammatory Cytokine Levels in Long-term OVX Rats Exposed to CUMS

CUMS significantly increased NF- κ B/p65/p50 levels in the hippocampus of SHAM rats compared to the non-CUMS control female rats ($P < 0.05$, (Fig. 7A)). CUMS resulted in a decrease of NF- κ B/p50/p65 contents in long-term OVX rats

compared to non-CUMS/CUMS SHAM rats ($F(5,12) = 28.44$ and $F(5,12) = 26.04$, $P < 0.05$, (Fig. 7A, B)). VD₃ (5.0 mg/kg, s.c.) supplementation reversed NF-kB/p50/p65 levels in long-term OVX/CUMS rats compared to the OVX/CUMS/solvent rats ($P < 0.05$, (Fig. 7A, B)). No significant differences of fluoxetine or paroxetine were found on hippocampal NF-kB/p50/p65 levels of the long-term OVX rats exposed to

CUMS when compared to the long-term OVX with CUMS rats treated with solvent ($P > 0.05$; (Fig. 7A, B)). The combination of SSRIs with VD₃ significantly decreased NF-kB/p50/p65 concentrations in the hippocampus in long-term OVX/CUMS rats when compared to the OVX/CUMS/solvent rats ($P < 0.05$, (Fig. 7A, B)).






Fig. (7). Effects of combined treatment with SSRIs plus VD₃ modulates hippocampal NF-κB/p50/p65 (A,B) and pro-inflammatory cytokines (C-E) levels in long-term OVX rats exposed to CUMS. * $P < 0.05$ versus the control group, # $P < 0.05$ versus the SHAM group exposed to CUMS, \$ $P < 0.05$ versus the OVX group subjected to CUMS, \$\$ $P < 0.05$ versus the OVX group subjected to CUMS treated with fluoxetine or paroxetine, respectively. The data are presented as mean \pm SD; $n = 7$ in each group.

TNF- α , IL-1 β , IL-6 concentrations in the hippocampus of SHAM rats were significantly increased compared to the non-CUMS control female rats ($P < 0.05$, (Fig 7C-E)). CUMS elevated hippocampal TNF- α , IL-1 β , IL-6 levels in the long-term OVX rats compared to the non-CUMS/CUMS SHAM rats ($F(5,12) = 28.44$, ($F(5,12) = 16.13$, ($F(5,12) = 22.10$, $P < 0.05$, (Fig. 7C-E)). VD₃ (5.0 mg/kg, s.c.) restored TNF- α , IL-1 β , IL-6 levels in long-term OVX/CUMS rats compared to the OVX/CUMS/solvent rats ($P < 0.05$, (Fig. 7C-E)). Fluoxetine and paroxetine induced a decrease of TNF- α , IL-1 β , IL-6 concentrations in long-term OVX/CUMS rats compared to the OVX/CUMS/solvent rats ($P < 0.05$, (Fig. 7B-D)). Co-treatment with SSRIs along with VD₃ completely restored TNF- α , IL-1 β , IL-6 levels in long-term OVX rats when compared with OVX/CUMS/fluoxetine/paroxetine/solvent and SHAM/CUMS/solvent rats ($P < 0.05$, (Fig. 7C-E)).

3.7. Combined Treatment (SSRIs with VD₃) Restores Hippocampal 5-HT and 5-HIAA Levels in Long-term OVX rats Exposed to CUMS

The HPLC assay showed that CUMS decreased 5-HT levels in the hippocampus of SHAM rats as compared to non-

CUMS control female rats ($P < 0.05$, (Fig. 8A)). In addition, the SHAM rats with CUMS showed a significantly enhanced 5-HT turnover in the hippocampus as compared to the non-CUMS control group. The post-hoc test indicated that long-term OVX rats with CUMS showed a more marked decrease of 5-HT concentrations in the hippocampus as compared to the non-CUMS/SHAM/CUMS rats ($F(5,12) = 22.84$, $P < 0.05$, (Fig. 8A)). A significant increase in 5-HIAA contents was detected in the hippocampus of long-term OVX rats with CUMS as compared to the non-CUMS/SHAM/CUMS rats ($F(5,12) = 34.56$, $P < 0.05$, (Fig. 8B)).

VD₃ treatment (5.0 mg/kg, s.c.) restored 5-HT and 5-HIAA levels of OVX/CUMS rats as compared to the OVX/CUMS/solvent group ($P < 0.05$, (Fig. 8A, B)). Treatment with SSRIs alone significantly elevated 5-HT and 5-HIAA levels in long-term OVX/CUMS rats as compared to the OVX/CUMS/solvent and SHAM/CUMS rats ($P < 0.05$, (Fig. 8A, B)). SSRIs in a combination with VD₃ resulted in normalization of 5-HT and 5-HIAA levels in long-term OVX rats as compared to the OVX/CUMS/fluoxetine/ paroxetine/ solvent and SHAM/CUMS/solvent rats ($P < 0.05$, (Fig. 8A, B)).




Fig. (8). Combined treatment with SSRIs plus VD₃ changes hippocampal 5-HT and 5-HIAA levels (A,B) in long-term OVX rats exposed to CUMS. * P < 0.05 versus the control group, # – P < 0.05 versus the SHAM group exposed to CUMS, \$ – P < 0.05 versus the OVX group subjected to CUMS, \$\$ – P < 0.05 versus the OVX group subjected to CUMS treated with fluoxetine or paroxetine, respectively. The data are presented as mean ± SD; n = 7 in each group.

4. DISCUSSION

This study aimed to examine the antidepressant-like effects of combined treatment of different SSRIs (fluoxetine and paroxetine) along with VD₃. A variety of behavioral tests were conducted on adult female rats with long-term lack of estrogen subjected to CUMS. The potential mechanism underlying the beneficial effects of a combination of SSRIs along with VD₃ was examined. To examine if this combination of drugs could effectively diminish the behavioral impairments produced by CUMS, we explored proinflammatory cytokines, NF-kB/p50/p65, and 5-HT/5-HIAA levels in the hippocampus. For the first time, proinflammatory cytokines, NF-kB/p50/p65, corticosterone/estradiol, and 5-HT/5-HIAA levels in the hippocampus were studied to evaluate the cooperative action of SSRIs and VD₃ in adult long-term OVX rats exposed to CUMS.

The CUMS model is regarded as one of the most frequently experimented models of depression and anhedonia [47, 48]. Laboratory animals subjected to CUMS show patterns of

behavioral alterations that are considered as experimental correlates of depression in humans [54, 55]. After being subjected to the CUMS protocols, SPT and FST were used to examine the anhedonia profile and behavioral despair in female rats, respectively. SPT is a standard test for the anhedonia-like state, which is postulated as a core sign of depression-like behavior in experimental models of depression [55]. FST is one of the most commonly used animal tests to assess the depression-like state and antidepressant-like effects of drugs in animals [56]. In this study, we examined the sucrose preference and body weight every week for 7 weeks to control the efficacy of CUMS protocol and concluded that 6 weeks of CUMS trials induced behavioral alterations in all experimental groups.

The results of this study show that SHAM/CUMS rats displayed decreased body weight, high anhedonia, depression-like, and anxiety-like symptoms in the SPT, FST, and OFT tests when compared with control non-CUMS animals. Moreover, SHAM/CUMS rats demonstrated elevated serum CS levels and hippocampal NF-kB/p50/p65, TNF-a, IL-1β, IL-6

levels, and reduced E_2 and 5-HT/5-HIAA concentrations in the hippocampus as compared to the control non-CUMS animals.

As compared to SHAM/CUMS rats, the decrease in body weight, as well as symptoms of anhedonia and depression in adult long-term OVX rats were more obvious when assessed by SPT and FST, respectively. Long-term OVX rats exposed to CUMS exhibited decreased locomotor and rearing activities and increased anxiety-like behavior in the OFT. Our data agree with other findings indicating that long-term estrogen deprivation in female rodents subjected to a stress-related procedure results in a profound depression-like profile [57]. This study's results demonstrated higher CS, NF- κ B/p50/p65, TNF- α , IL-1 β , and IL-6 levels in adult long-term OVX rats subjected to CUMS as compared to the SHAM/CUMS rats. Estrogen deficiency contributes to the impaired production of inflammatory cytokines in the whole organism [58]. These alterations in dysregulation in the production of inflammatory cytokines may extend to the brain and induce a state of neuroinflammation in the brain, which is directly associated with the development of affective-related disorders [15 - 21]. Our data are in an agreement with the experimental findings showing that ovariectomy elevates the production of proinflammatory cytokines [58]. The long-term OVX rats with CUMS demonstrated a decrease in 5-HT/5-HIAA levels in the hippocampus as compared to SHAM/CUMS rats. The current results confirm that CUMS produced marked behavioral, neuroendocrine, and neuroimmunological changes in adult SHAM or OVX rats with long-term estrogen deprivation (post-ovariectomy period of three months). These findings agree with our previous data and data reported in other studies [54, 55, 59].

After 28 days of therapy, we found that VD₃ (5.0 mg/kg, s.c.) decreased the anhedonia and depression-like states in OVX/CUMS rats as published previously [42]. Moreover, decreased serum CS and hippocampal NF- κ B/p50/p65, TNF- α , IL-1 β , IL-6 levels, as well as increased E_2 and 5-HT/5-HIAA concentrations in the hippocampus were also noted in OVX/CUMS animals treated with VD₃ for 4 weeks as compared to the OVX/CUMS rats.

Treatment with the standard antidepressants, fluoxetine or paroxetine alone, induced a reduction of anhedonia and depression-like states in the SPT and FST from the 5th week of treatment, along with a decrease in the anxiety-like profile in the OFT of long-term OVX/CUMS rats as compared to OVX/SHAM/CUMS/solvent females. Simultaneously, long-term OVX/CUMS rats administered with tested SSRIs demonstrated a decrease of serum CS and hippocampal pro-inflammatory cytokine contents, and an increase of 5-HT/5-HIAA levels in the hippocampus as compared to OVX/SHAM/CUMS/solvent rats. However, fluoxetine and paroxetine did not modify NF- κ B/p50/p65 levels in the hippocampus of long-term OVX/CUMS rats submitted to CUMS. Our experimental data confirm that in a similar manner to other SSRIs, fluoxetine and paroxetine restore the functional activity of the HPA axis, 5-HT neurotransmission, and neuroinflammation in the brain, improving the depression-like state of OVX rats. Moreover, the data of the present study are in line with previous findings showing that paroxetine might

induce an increase of E_2 levels in females [45, 46].

Supplementation of VD₃ to fluoxetine treatment failed to enhance the antidepressant-like effects of both drugs in long-term OVX rats with CUMS as compared to OVX/CUMS females treated with fluoxetine alone. However, this combination produced a profound decrease in serum CS concentrations and an increase of hippocampal 5-HT/5-HIAA levels in long-term OVX/CUMS rats in contrast to OVX/CUMS/fluoxetine rats. Moreover, only the application of fluoxetine with VD₃ increased E_2 levels and decreased NF- κ B/p50/p65 contents in long-term OVX/CUMS rats. A better comprehension of NF- κ B-dependent mechanisms in the antidepressant action of SSRIs along VD₃ requires further research. Co-administration of paroxetine with VD₃ quickly and more effectively produced antianhedonic- and antidepressant-like effects in long-term OVX rats with CUMS as compared to treatment with paroxetine alone. In contrast to the behavioral effects of fluoxetine and VD₃, long-term OVX/CUMS rats after 1 week of treatment with paroxetine and VD₃ demonstrated a marked decrease in anhedonia and depression-like symptoms. Furthermore, co-treatment with paroxetine and VD₃ produced synergic effects in decreasing serum CS and E_2 levels as well as hippocampal pro-inflammatory cytokines contents.

In addition, enhancement of 5-HT/5-HIAA concentrations in the hippocampus was registered in long-term OVX female rats exposed to CUMS.

Interestingly, fluoxetine or paroxetine administered with VD₃ cooperatively increased the total locomotor activity, rearings, the number of crossings, and time spent in central squares in the OFT in long-term OVX/CUMS rats as compared to in OVX/CUMS rats treated with fluoxetine or paroxetine alone. The number of crossings and time spent in central squares in the OFT are often interpreted as indicators of fear or anxiety in laboratory animals. Increases in these parameters can be induced by diazepam and other anxiolytics [60, 61]. All of the used drugs in the current study increased the number of crossings and time spent in central squares in the OFT in long-term OVX/CUMS rats. These results led us to conclude that fluoxetine or paroxetine administered with VD₃ might similarly affect the anxiety-like profile in long-term OVX/CUMS rats as compared to its different effects on depression-like behavior in the FST. Based on these data, our next study will aim to assess exploratory activity and anxiety-like behavior in long-term OVX rats exposed to CUMS treated with fluoxetine and paroxetine along with VD₃ using more specific tests for anxiety.

In conclusion, the synergistic effect of co-administration of paroxetine (10.0 mg/kg, i.p.) and VD₃ (5.0 mg/kg, s.c.) completely abolished the CUMS-induced behavioral, neurohormonal, and neuroimmunological impairments in long-term OVX/CUMS rats. The supplementation of VD₃ to fluoxetine failed to enhance the efficacy of this SSRI in long-term OVX/CUMS rats. However, it is important to note that co-treatment with fluoxetine and VD₃ induces an increase in serum E_2 concentrations and NF- κ B/p50/p65 levels in the hippocampus of long-term OVX/CUMS rats. Therefore, it can

be inferred that the efficacy of a combination of SSRIs with VD₃ depends upon the SSRI used. This is why clinical and preclinical findings regarding the efficacy of a combination of SSRIs with VD₃ are highly debated. [31, 62].

VD₃ deficiency has been proven to impact the pathogenesis of various diseases, including mood disorders [26 - 29]. Moreover, VD₃ is involved in the control of NF-κB signaling, 5-HT neurotransmission, and neuroinflammation in the brain [32 - 35]. Low VD₃ levels are found in the majority of postmenopausal women [36 - 38]. Therefore, VD₃ supplementation may be useful in treating mood disturbances in postmenopausal women with a low level of VD₃. However, the exact role of VD₃ supplementation in the prevention and treatment of mood disorders associated with menopausal consequences has not been completely identified. On the other hand, estrogen deficiency, like VD₃ deficiency, also play a part in the mechanisms of affective-related disorders by modulating 5-HT system and pro-inflammatory cytokine levels in the brain [26, 27]. Such a complex interaction between estrogen and VD₃ creates a neurobiological basis for their participation in similar behavioral disorders. Moreover, several investigations on the effects of VD₃ on the brain have documented that VD₃ can alter the synthesis of steroid hormones, including estrogen [63 - 66].

NF-κB is activated by different trigger factors stimulated by neuroinflammation [24, 25]. The up-regulation of NF-κB activity has been found in the experimental stress model [24, 25]. Pro-inflammatory cytokines, including IL-1β, IL-6, and TNF-α, are strongly involved in the pathophysiology of depression [17 - 20]. Some reports showed increased pro-inflammatory cytokine levels in patients with depression [18 - 21]. On the other hand, pro-inflammatory cytokines modulate the monoamines' functional activity and turnover of 5-HT in the brain [67 - 70]. Therefore, we believe that the beneficial effects of paroxetine and VD₃ in the long-term OVX/CUMS rats are associated with the cooperative effects of these drugs on serum CS and E₂ levels, and hippocampal TNF-α, IL-1β, IL-6, and 5-HT/5-HIA concentrations.

Most of the current SSRIs exert an antidepressant-like effect by enhancing the 5-HT and/or other monoamines levels [5 - 9]. 5-HT is the main neurotransmitter that controls normal hippocampal neurogenesis [6, 7]. Therefore, we can assume that the antidepressant-like activity of SSRIs along with VD₃ in long-term OVX rats subjected to CUMS is associated with complex positive effects of paroxetine and VD₃ on depression-like behavior in CUMS model *via* restoration of 5-HT, as well as NF-κB/p50/p65 and pro-inflammatory cytokine levels in the brain. Further research is needed to clarify this hypothesis.

Among different classes of antidepressants, SSRIs are the most extensively used pharmacological drugs to treat affective-related profiles in perimenopausal and postmenopausal women [1 - 6]. The combined application of drugs is often used to achieve more considerable and faster therapeutic responses in the treatment of depression [71]. Therefore, VD₃ supplementation to SSRIs may be effective in treating affective-related disturbances among hypoestrogenic females with VD₃ deficiencies or insufficiencies. However, more research on this topic is still required since the exact role of VD₃ in affective-

related disorders associated with hormonal changes in women has not been completely understood.

The findings of this study provide experimental evidence regarding the beneficial effects of co-treatment with paroxetine along with VD₃ in a CUMS model, producing cooperative antianhedonic- and anti-depressant-like effects in the long-term OVX adult rats. Further research should explore the precise mechanism behind paroxetine and VD₃ co-treatment in long-term OVX rats subjected to CUMS.

CONCLUSION

In conclusion, the results of this study suggest that combined treatment with fluoxetine (10.0 mg/kg, i.p.) or paroxetine (10.0 mg/kg, i.p.) along with VD₃ (5.0 mg/kg, s.c.) produces distinct effects on depression-like behavior in long-term OVX rats exposed to CUMS. Co-administration of fluoxetine (10.0 mg/kg, i.p.) along with VD₃ (5.0 mg/kg, s.c.) did not facilitate the effectiveness of fluoxetine in long-term OVX rats submitted to CUMS. The application of paroxetine (10.0 mg/kg, i.p.) along with VD₃ (5.0 mg/kg, s.c.) releases faster and more considerable antianhedonic- and anti-depressant-like effects in long-term OVX rats with CUMS than treatment with paroxetine alone. Our data indicate that the co-administration of paroxetine and VD₃ may attenuate stress-induced CS/E₂ and proinflammatory cytokine/NF-κB/5-HT alterations in the hippocampus of long-term OVX rats exposed to CUMS. This is the first report of the application of these drugs in combination to relieve anhedonia and depression-like state in long-term OVX rats exposed to CUMS. Our findings suggest that paroxetine and VD₃ may be a therapeutic alternative to ameliorate the depression associated with menopause in women who have VD₃ deficiency. This work provides translational information for the development of effective therapies for menopausal mood disorders in the future.

ETHICAL APPROVAL AND CONSENT TO PARTICIPATE

The study is approved by the Animal Care Committee of the I.P. Pavlov Institute of Physiology, Russian Academy of Sciences, Russian Federation, protocol 1095/1/25.06.2012.

HUMAN AND ANIMAL RIGHTS

No humans were used in this study. The current experiments on female rats were in accordance with the guidelines of National Institute of Health (National Research Council, Publication No. 85-23, revised in 1996), and the Animal Welfare Assurance Renewal for the I.P. Pavlov Institute of Physiology.

CONSENT FOR PUBLICATION

Not applicable.

AVAILABILITY OF DATA AND MATERIALS

The data supporting the findings of this article is available at <https://grant.rscf.ru> upon the request.

FUNDING

The present study was supported by the Russian Science Foundation (RSF), research project No. 16-15-10053 (extension).

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Frey BN, Lord C, Soares CN. Depression during menopausal transition: A review of treatment strategies and pathophysiological correlates. *Menopause Int* 2008; 14(3): 123-8. [http://dx.doi.org/10.1258/mi.2008.008019] [PMID: 18714078]
- [2] Luppá M, Sikorski C, Luck T, *et al.* Age- and gender-specific prevalence of depression in latest-life: Systematic review and meta-analysis. *J Affect Disord* 2012; 136(3): 212-21. [http://dx.doi.org/10.1016/j.jad.2010.11.033] [PMID: 21194754]
- [3] Garcia-Portilla MP. Depression and perimenopause: A review. *Actas Esp Psiquiatr* 2009; 37(4): 213-21. [PMID: 19927235]
- [4] Craske MG, Stein MB. Anxiety. *Lancet* 2016; 388(10063): 3048-59. [http://dx.doi.org/10.1016/S0140-6736(16)30381-6] [PMID: 27349358]
- [5] Heidari M, Ghodusi M, Rafiei H. Sexual self-concept and its relationship to depression, stress and anxiety in postmenopausal women. *J Menopausal Med* 2017; 23(1): 42-8. [http://dx.doi.org/10.6118/jmm.2017.23.1.42] [PMID: 28523258]
- [6] Sansone RA, Sansone LA. Antidepressant adherence: Are patients taking their medications? *Innov Clin Neurosci* 2012; 9(5-6): 41-6. [PMID: 22808448]
- [7] Cipriani A, Furukawa TA, Salanti G, *et al.* Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: A systematic review and network meta-analysis. *Lancet* 2018; 391(10128): 1357-66. [http://dx.doi.org/10.1016/S0140-6736(17)32802-7] [PMID: 29477251]
- [8] Arroll B, Macgillivray S, Ogston S, *et al.* Efficacy and tolerability of tricyclic antidepressants and SSRIs compared with placebo for treatment of depression in primary care: A meta-analysis. *Ann Fam Med* 2005; 3(5): 449-56. [http://dx.doi.org/10.1370/afm.349] [PMID: 16189062]
- [9] Cassano P, Fava M. Tolerability issues during long-term treatment with antidepressants. *Ann Clin Psychiatry* 2004; 16(1): 15-25. [http://dx.doi.org/10.1080/10401230490281618] [PMID: 15147109]
- [10] Dording CM, Mischoulon D, Petersen TJ, *et al.* The pharmacologic management of SSRI-induced side effects: A survey of psychiatrists. *Ann Clin Psychiatry* 2002; 14(3): 143-7. [http://dx.doi.org/10.3109/10401230209147450] [PMID: 12585563]
- [11] Anderson IM. Selective serotonin reuptake inhibitors *versus* tricyclic antidepressants: A meta-analysis of efficacy and tolerability *J Affect Disord* 2000; 58(1): 19-36. [http://dx.doi.org/10.1016/S0165-0327(99)00092-0] [PMID: 10760555]
- [12] Moncrieff J. Does antipsychotic withdrawal provoke psychosis? Review of the literature on rapid onset psychosis (supersensitivity psychosis) and withdrawal-related relapse. *Acta Psychiatr Scand* 2006; 114(1): 3-13. [http://dx.doi.org/10.1111/j.1600-0447.2006.00787.x] [PMID: 16774655]
- [13] Graziottin A, Serafini A. Depression and the menopause: Why antidepressants are not enough? *Menopause Int* 2009; 15(2): 76-81. [http://dx.doi.org/10.1258/mi.2009.009021] [PMID: 19465674]
- [14] Kraus C, Castrén E, Kasper S, Lanzenberger R. Serotonin and neuroplasticity - Links between molecular, functional and structural pathophysiology in depression. *Neurosci Biobehav Rev* 2017; 77: 317-26. [http://dx.doi.org/10.1016/j.neubiorev.2017.03.007] [PMID: 28342763]
- [15] Miller AH, Maletic V, Raison CL. Inflammation and its discontents: The role of cytokines in the pathophysiology of major depression. *Biol Psychiatry* 2009; 65(9): 732-41. [http://dx.doi.org/10.1016/j.biopsych.2008.11.029] [PMID: 19150053]
- [16] Rosenblat JD, Cha DS, Mansur RB, McIntyre RS. Inflamed moods: A review of the interactions between inflammation and mood disorders. *Prog Neuropsychopharmacol Biol Psychiatry* 2014; 53: 23-34. [http://dx.doi.org/10.1016/j.pnpbp.2014.01.013] [PMID: 24468642]
- [17] You Z, Luo C, Zhang W, *et al.* Pro- and anti-inflammatory cytokines expression in rat's brain and spleen exposed to chronic mild stress: Involvement in depression. *Behav Brain Res* 2011; 225(1): 135-41. [http://dx.doi.org/10.1016/j.bbr.2011.07.006] [PMID: 21767575]
- [18] Liu Y, Ho RC, Mak A. Interleukin (IL)-6, tumour necrosis factor alpha (TNF- α) and soluble interleukin-2 receptors (sIL-2R) are elevated in patients with major depressive disorder: A meta-analysis and meta-regression. *J Affect Disord* 2012; 139(3): 230-9. [http://dx.doi.org/10.1016/j.jad.2011.08.003] [PMID: 21872339]
- [19] Sukoff Rizzo SJ, Neal SJ, Hughes ZA, *et al.* Evidence for sustained elevation of IL-6 in the CNS as a key contributor of depressive-like phenotypes. *Transl Psychiatry* 2012; 2: e199. [http://dx.doi.org/10.1038/tp.2012.120] [PMID: 23212583]
- [20] Réus GZ, Fries GR, Stertz L, *et al.* The role of inflammation and microglial activation in the pathophysiology of psychiatric disorders. *Neuroscience* 2015; 300: 141-54. [http://dx.doi.org/10.1016/j.neuroscience.2015.05.018] [PMID: 25981208]
- [21] Dunn AJ, Swiergiel AH, de Beaupaire R. Cytokines as mediators of depression: What can we learn from animal studies? *Neurosci Biobehav Rev* 2005; 29(4-5): 891-909. [http://dx.doi.org/10.1016/j.neubiorev.2005.03.023] [PMID: 15885777]
- [22] McKay LI, Cidlowski JA. Molecular control of immune/inflammatory responses: Interactions between nuclear factor-kappa B and steroid receptor-signaling pathways. *Endocr Rev* 1999; 20(4): 435-59. [PMID: 10453354]
- [23] Caviedes A, Lafourcade C, Soto C, Wyneken U. BDNF/NF- κ B signaling in the neurobiology of depression. *Curr Pharm Des* 2017; 23(21): 3154-63. [http://dx.doi.org/10.2174/1381612823666170111141915] [PMID: 28078988]
- [24] Lindström TM, Bennett PR. The role of nuclear factor kappa B in human labour. *Reproduction* 2005; 130(5): 569-81. [http://dx.doi.org/10.1530/rep.1.00197] [PMID: 16264088]
- [25] Mohamed MR, McFadden G. NF κ B inhibitors: Strategies from poxviruses. *Cell Cycle* 2009; 8(19): 3125-32. [http://dx.doi.org/10.4161/cc.8.19.9683] [PMID: 19738427]
- [26] Villa A, Vegeto E, Poletti A, Maggi A. Estrogens, neuroinflammation, and neurodegeneration. *Endocr Rev* 2016; 37(4): 372-402. [http://dx.doi.org/10.1210/er.2016-1007] [PMID: 27196727]
- [27] Amin Z, Canli T, Epperson CN. Effect of estrogen-serotonin interactions on mood and cognition. *Behav Cogn Neurosci Rev* 2005; 4(1): 43-58. [http://dx.doi.org/10.1177/1534582305277152] [PMID: 15886402]
- [28] Spedding S. Vitamin D and depression: A systematic review and meta-analysis comparing studies with and without biological flaws. *Nutrients* 2014; 6(4): 1501-18. [http://dx.doi.org/10.3390/nu6041501] [PMID: 24732019]
- [29] Chapuy MC, Preziosi P, Maamer M, *et al.* Prevalence of vitamin D insufficiency in an adult normal population. *Osteoporos Int* 1997; 7(5): 439-43. [http://dx.doi.org/10.1007/s001980050030] [PMID: 9425501]
- [30] Holick MF, Chen TC. Vitamin D deficiency: A worldwide problem with health consequences. *Am J Clin Nutr* 2008; 87(4): 1080S-6S. [http://dx.doi.org/10.1093/ajcn/87.4.1080S] [PMID: 18400738]
- [31] Bertone-Johnson ER, Powers SI, Spangler L, *et al.* Vitamin D intake from foods and supplements and depressive symptoms in a diverse population of older women. *Am J Clin Nutr* 2011; 94(4): 1104-12. [http://dx.doi.org/10.3945/ajcn.111.017384] [PMID: 21865327]
- [32] Kesby JP, Eyles DW, Burne TH, McGrath JJ. The effects of vitamin D on brain development and adult brain function. *Mol Cell Endocrinol* 2011; 347(1-2): 121-7. [http://dx.doi.org/10.1016/j.mce.2011.05.014] [PMID: 21664231]
- [33] Eyles DW, Burne TH, McGrath JJ. Vitamin D, effects on brain development, adult brain function and the links between low levels of vitamin D and neuropsychiatric disease. *Front Neuroendocrinol* 2013; 34(1): 47-64.

- [34] <http://dx.doi.org/10.1016/j.yfrne.2012.07.001> [PMID: 22796576]
Garcion E, Wion-Barbot N, Montero-Menei CN, Berger F, Wion D. New clues about vitamin D functions in the nervous system. *Trends Endocrinol Metab* 2002; 13(3): 100-5.
[http://dx.doi.org/10.1016/S1043-2760\(01\)00547-1](http://dx.doi.org/10.1016/S1043-2760(01)00547-1) [PMID: 11893522]
- [35] Norlin M. Effects of vitamin D in the nervous system: Special focus on interaction with steroid hormone signalling and a possible role in the treatment of brain cancer. *J Neuroendocrinol* 2020; 32(1): e12799.
<http://dx.doi.org/10.1111/jne.12799> [PMID: 31593305]
- [36] LeBlanc ES, Desai M, Perrin N, *et al.* Vitamin D levels and menopause-related symptoms. *Menopause* 2014; 21(11): 1197-203.
<http://dx.doi.org/10.1097/GME.000000000000238> [PMID: 24736200]
- [37] Milanesechi Y, Shardell M, Corsi AM, *et al.* Serum 25-hydroxyvitamin D and depressive symptoms in older women and men. *J Clin Endocrinol Metab* 2010; 95(7): 3225-33.
<http://dx.doi.org/10.1210/jc.2010-0347> [PMID: 20444911]
- [38] Parker GB, Brotchie H, Graham RK. Vitamin D and depression. *J Affect Disord* 2017; 208: 56-61.
<http://dx.doi.org/10.1016/j.jad.2016.08.082> [PMID: 27750060]
- [39] Sarris J, Murphy J, Mischoulon D, *et al.* Adjunctive nutraceuticals for depression: A systematic review and meta-analyses. *Am J Psychiatry* 2016; 173(6): 575-87.
<http://dx.doi.org/10.1176/appi.ajp.2016.15091228> [PMID: 27113121]
- [40] Delvin E, Souberbielle JC, Viard JP, Salle B. Role of vitamin D in acquired immune and autoimmune diseases. *Crit Rev Clin Lab Sci* 2014; 51(4): 232-47.
<http://dx.doi.org/10.3109/10408363.2014.901291> [PMID: 24813330]
- [41] Peelen E, Knippenberg S, Muris A-H, *et al.* Effects of vitamin D on the peripheral adaptive immune system: A review. *Autoimmun Rev* 2011; 10(12): 733-43.
<http://dx.doi.org/10.1016/j.autrev.2011.05.002> [PMID: 21621002]
- [42] Koshkina A, Dudnichenko T, Baranenko D, Fedotova J, Drago F. Effects of Vitamin D3 in long-term ovariectomized rats subjected to chronic unpredictable mild stress: GDNF, NT-3, and NT-4 implications. *Nutrients* 2019; 11(8): E1726.
<http://dx.doi.org/10.3390/nu11081726> [PMID: 31357443]
- [43] National Institute for Clinical Excellence. Depression: Management of depression in primary and secondary care 2004. www.nice.org.uk
- [44] Magni LR, Purgato M, Gastaldon C, *et al.* Fluoxetine versus other types of pharmacotherapy for depression *Cochrane Database Syst Rev* 2013; 7(7): CD004185.
<http://dx.doi.org/10.1002/14651858.CD004185.pub3> [PMID: 24353997]
- [45] Cubeddu A, Giannini A, Bucci F, *et al.* Paroxetine increases brain-derived neurotrophic factor in postmenopausal women. *Menopause* 2010; 17(2): 338-43.
<http://dx.doi.org/10.1097/gme.0b013e3181c29e44> [PMID: 19934779]
- [46] Soares CN, Joffe H, Viguera AC, *et al.* Paroxetine versus placebo for women in midlife after hormone therapy discontinuation *Am J Med* 2008; 121(2): 159-162.e1.
<http://dx.doi.org/10.1016/j.amjmed.2007.10.007> [PMID: 18261506]
- [47] Katz RJ. Animal models and human depressive disorders. *Neurosci Biobehav Rev* 1981; 5(2): 231-46.
[http://dx.doi.org/10.1016/0149-7634\(81\)90004-X](http://dx.doi.org/10.1016/0149-7634(81)90004-X) [PMID: 7022272]
- [48] Willner P, Towell A, Sampson D, Sophokleous S, Muscat R. Reduction of sucrose preference by chronic unpredictable mild stress, and its restoration by a tricyclic antidepressant. *Psychopharmacology (Berl)* 1987; 93(3): 358-64.
<http://dx.doi.org/10.1007/BF00187257> [PMID: 3124165]
- [49] Anisman H, Matheson K. Stress, depression, and anhedonia: Caveats concerning animal models. *Neurosci Biobehav Rev* 2005; 29(4-5): 525-46.
<http://dx.doi.org/10.1016/j.neubiorev.2005.03.007> [PMID: 15925696]
- [50] Amodeo LR, Greenfield VY, Humphrey DE, *et al.* Effects of acute or repeated paroxetine and fluoxetine treatment on affective behavior in male and female adolescent rats. *Psychopharmacology (Berl)* 2015; 232(19): 3515-28.
<http://dx.doi.org/10.1007/s00213-015-4003-1> [PMID: 26141193]
- [51] Heffner TG, Hartman JA, Seiden LS. A rapid method for the regional dissection of the rat brain. *Pharmacol Biochem Behav* 1980; 13(3): 453-6.
[http://dx.doi.org/10.1016/0091-3057\(80\)90254-3](http://dx.doi.org/10.1016/0091-3057(80)90254-3) [PMID: 7422701]
- [52] Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976; 72: 248-54.
[http://dx.doi.org/10.1016/0003-2697\(76\)90527-3](http://dx.doi.org/10.1016/0003-2697(76)90527-3) [PMID: 942051]
- [53] Fedotova J, Zarembo D, Dragasek J, Caprnda M, Kruzliak P, Dudnichenko T. Modulating effects of cholecalciferol treatment on estrogen deficiency-induced anxiety-like behavior of adult female rats. *Folia Med (Plovdiv)* 2017; 59(2): 139-58.
<http://dx.doi.org/10.1515/folmed-2017-0022> [PMID: 28704190]
- [54] Burstein O, Franko M, Gale E, *et al.* Escitalopram and NHT normalized stress-induced anhedonia and molecular neuroadaptations in a mouse model of depression. *PLoS One* 2017; 12(11): e0188043.
<http://dx.doi.org/10.1371/journal.pone.0188043> [PMID: 29141007]
- [55] Matthews K, Forbes N, Reid IC. Sucrose consumption as an hedonic measure following chronic unpredictable mild stress. *Physiol Behav* 1995; 57(2): 241-8.
[http://dx.doi.org/10.1016/0031-9384\(94\)00286-E](http://dx.doi.org/10.1016/0031-9384(94)00286-E) [PMID: 7716198]
- [56] Huang H, Zhao J, Jiang L, *et al.* Paconiflorin improves menopause depression in ovariectomized rats under chronic unpredictable mild stress. *Int J Clin Exp Med* 2015; 8(4): 5103-11.
[PMID: 26131083]
- [57] Sedaghat K, Yousefian Z, Vafaei AA, *et al.* Mesolimbic dopamine system and its modulation by vitamin D in a chronic mild stress model of depression in the rat. *Behav Brain Res* 2019; 356: 156-69.
<http://dx.doi.org/10.1016/j.bbr.2018.08.020> [PMID: 30144460]
- [58] Abu-Taha M, Rius C, Hermenegildo C, *et al.* Menopause and ovariectomy cause a low grade of systemic inflammation that may be prevented by chronic treatment with low doses of estrogen or losartan. *J Immunol* 2009; 183(2): 1393-402.
<http://dx.doi.org/10.4049/jimmunol.0803157> [PMID: 19553526]
- [59] Jiang H, Wang Z, Wang Y, *et al.* Antidepressant-like effects of curcumin in chronic mild stress of rats: Involvement of its anti-inflammatory action. *Prog Neuropsychopharmacol Biol Psychiatry* 2013; 47: 33-9.
<http://dx.doi.org/10.1016/j.pnpbp.2013.07.009> [PMID: 23876788]
- [60] Shaw D, Annett JM, Doherty B, Leslie JC. Anxiolytic effects of lavender oil inhalation on open-field behaviour in rats. *Phytomedicine* 2007; 14(9): 613-20.
<http://dx.doi.org/10.1016/j.phymed.2007.03.007> [PMID: 17482442]
- [61] Hata T, Nishimura Y, Kita T, Itoh E, Kawabata A. The abnormal open-field behavior of SART-stressed rats and effects of some drugs on it. *Jpn J Pharmacol* 1988; 48(4): 479-90.
<http://dx.doi.org/10.1254/jjp.48.479> [PMID: 3244201]
- [62] Gaugris S, Heaney RP, Boonen S, Kurth H, Bentkover JD, Sen SS. Vitamin D inadequacy among post-menopausal women: A systemic review. *QJM: Month J Assoc Physicians* 2005; 98: 667-76.
- [63] Bakhshalizadeh S, Amidi F, Shirazi R, Shabani Nashtaei M. Vitamin D3 regulates steroidogenesis in granulosa cells through AMP-activated protein kinase (AMPK) activation in a mouse model of polycystic ovary syndrome. *Cell Biochem Funct* 2018; 36(4): 183-93.
<http://dx.doi.org/10.1002/cbf.3330> [PMID: 29676471]
- [64] Lundqvist J, Norlin M, Wikvall K. 1alpha,25-Dihydroxyvitamin D3 affects hormone production and expression of steroidogenic enzymes in human adrenocortical NCI-H295R cells. *Biochim Biophys Acta* 2010; 1801(9): 1056-62.
<http://dx.doi.org/10.1016/j.bbali.2010.04.009> [PMID: 20420936]
- [65] Lundqvist J, Norlin M, Wikvall K. 1alpha,25-Dihydroxyvitamin D3 exerts tissue-specific effects on estrogen and androgen metabolism. *Biochim Biophys Acta* 2011; 1811(4): 263-70.
<http://dx.doi.org/10.1016/j.bbali.2011.01.004> [PMID: 21262387]
- [66] Yague JG, Garcia-Segura LM, Azcoitia I. Selective transcriptional regulation of aromatase gene by vitamin D, dexamethasone, and mifepristone in human glioma cells. *Endocrine* 2009; 35(2): 252-61.
<http://dx.doi.org/10.1007/s12020-008-9134-2> [PMID: 19116788]
- [67] Maes M. Evidence for an immune response in major depression: A review and hypothesis. *Prog Neuropsychopharmacol Biol Psychiatry* 1995; 19(1): 11-38.
[http://dx.doi.org/10.1016/0278-5846\(94\)00101-M](http://dx.doi.org/10.1016/0278-5846(94)00101-M) [PMID: 7708925]
- [68] Zhu CB, Blakely RD, Hewlett WA. The proinflammatory cytokines interleukin-1beta and tumor necrosis factor-alpha activate serotonin transporters. *Neuropsychopharmacology* 2006; 31(10): 2121-31.
<http://dx.doi.org/10.1038/sj.npp.1301029> [PMID: 16452991]
- [69] Kong E, Susic S, Monje FJ, *et al.* STAT3 controls IL6-dependent regulation of serotonin transporter function and depression-like behavior. *Sci Rep* 2015; 5: 9009.
<http://dx.doi.org/10.1038/srep09009> [PMID: 25760924]
- [70] Bryleva EY, Brundin L. Kynurenine pathway metabolites and

suicidality. *Neuropharmacology* 2017; 112(Pt B): 324-30.
[<http://dx.doi.org/10.1016/j.neuropharm.2016.01.034>]
26820800]

[PMID:

[71] Levinson AJ, Fitzgerald PB, Favalli G, Blumberger DM, Daigle M, Daskalakis ZJ. Evidence of cortical inhibitory deficits in major depressive disorder. *Biol Psychiatry* 2010; 67(5): 458-64.
[<http://dx.doi.org/10.1016/j.biopsych.2009.09.025>] [PMID: 19922906]

© 2020 Julia Fedotova.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.